



11. (New) A bathophenanthroline compound of formula (I):



wherein R^1 and R^2 are derived from R^1 -Li and R^2 -Li respectively, and may be the same or different and independently represent a hydrocarbon group provided that at least one of R^1 and R^2 has at least two carbons; and wherein R^1 and R^2 are selected from the group consisting of an ethyl group, an n-propyl group, an isopropyl group, an n-pentyl group, an iso-pentyl group, a neopentyl group, a tert-pentyl group, a cyclopentyl group, a methycyclopentyl group, a dimethycyclopentyl group, a trimethycyclopentyl group, a tetramethycyclopentyl group, an n-hexyl group, a 2-ethylbutyl group, a 3,3-dimethylbutyl group, a cyclohexyl group, an n-cyclohexylmethyl group, an n,n-dimethycyclohexyl group, an n,n,n-trimethycyclohexyl group, a tert-octyl group, a 2-ethylheptyl group, an n-nonyl group, an n-decyl group, an n-dodecyl group, an n-tetradecyl group, and an n-hexadecyl group, ~~wherein $n \geq 0$~~ a benzyl group, a phenethyl group, an α -methylbenzyl group, an α,α -dimethylbenzyl group, a 1-naphthylmethyl group, a 2-naphthylmethyl group, a furfuryl group, a 2-methylbenzyl group, a 3-methylbenzyl group, a 4-methylbenzyl group, a 4-ethylbenzyl group, a 4-isopropylbenzyl group, a 4-tert-butylbenzyl group, a 4-n-hexylbenzyl group, a 4-nonylbenzyl group, and a 3,4-dimethylbenzyl group.

12. Cancel.

13. (New) A bathophenanthroline compound adapted to be used as an organic layer having a luminescent region provided between an anode and a cathode, wherein the organic layer comprises a bathophenanthroline compound of formula (II):




wherein Ar^1 and Ar^2 may be the same or different and independently represent an aryl group but do not form an interlocking macrocyclic compound.

14. (New) The bathophenanthroline compound according to claim 13 wherein R^1 and R^2 Ar^1 and Ar^2 are selected from the group consisting of a 1-naphthyl group, a 2-anthryl group, a 9-anthryl group, a 2-fluorenyl group, a 4-quinolyl group, a pyridyl group, a 3-pyridynyl group, a 2-pyridynyl group, a 3-furyl group, a 2-furyl group, a 3-thienyl group, a 2-oxazolyl group, a 2-thiazolyl group, a 2-benzoxazolyl group, a 2-benzothiazolyl group, a 2-benzoimidazolyl group, a 4-methyphenyl group, a 3-methyphenyl group, a 2-methyphenyl group, a n,n-dimethyphenyl group, a n,n,n-trimethyphenyl group, a n-ethyphenyl group, a n,n-diethyphenyl group, a n,n,n-triethyphenyl group, a 4-n-propylphenyl group, a n-isopropylphenyl group, a 4-n-butylphenyl group, a 4-isobutylphenyl group, a 4-sec-butylphenyl group, a n-tert-butylphenyl group, ~~wherein~~ $n \geq 0$.


15. Cancel.

16. (New) A process, comprising:

(a) obtaining ~~subjecting~~ a bathophenanthroline of formula (IV)



(b) subjecting the bathophenanthroline to nucleophilic substitution reaction at the 2, 9 positions by a lithium compound of formula (III):




wherein R^1 and R^2 may be the same or different and independently represent a hydrocarbon group provided that at least one of R^1 and R^2 has at least two carbon atoms, and R^1 reacts at the 2 position and R^2 reacts at the 9 position of the bathophenanthroline.

17. (New) The process according to Claim 16, wherein a carbanion is generated from the lithium compound in a solution and reacted with the bathophenanthroline during the nucleophilic substitution reaction.


18. (New) A process, comprising:

(a) obtaining ~~subjecting~~ a lithium compound of formula (V):



wherein Ar^1 and Ar^2 may be the same or different and independently represent an aryl group,

(b) subjecting the lithium compound to a bathophenanthroline of formula (IV):



via nucleophilic substitution reaction at the 2, 9 positions of the bathophenanthroline where A^1 reacts at the 2 position and A^2 reacts at the 9 position of the bathophenanthroline.